

# PubMed

Search: wallach nik

U.S. National Library of Medicine  
National Institutes of Health

Filter your results: All (8)

Display Settings: Summary, 20 per page, Sorted by Recently Added

## Results: 8

1. NF-kappaB-inducing kinase is involved in the activation of the CD28 responsive element through phosphorylation of c-Rel and regulation of its transactivating activity.  
Sánchez-Valdepeñas C, Martin AG, Ramakrishnan P, Wallach D, Fresno M.  
J Immunol. 2006 Apr 15;176(8):4666-74.  
PMID: 16585559 [PubMed - indexed for MEDLINE]  
[Free article](#)
2. Receptor-specific signaling for both the alternative and the canonical NF-kappaB activation pathways by NF-kappaB-inducing kinase.  
Ramakrishnan P, Wang W, Wallach D.  
Immunity. 2004 Oct;21(4):477-89.  
PMID: 15495626 [PubMed - indexed for MEDLINE]
3. Induction of interleukin-8 synthesis integrates effects on transcription and mRNA degradation from at least three different cytokine- or stress-activated signal transduction pathways.  
Holtmann H, Winzen R, Holland P, Eickemeier S, Hoffmann E, Wallach D, Malinin NL, Cooper JA, Resch K, Kracht M.  
Mol Cell Biol. 1999 Oct;19(10):6742-53.  
PMID: 10490613 [PubMed - indexed for MEDLINE]  
[Free article](#)
4. Identification of a cell protein (FIP-3) as a modulator of NF-kappaB activity and as a target of an adenovirus inhibitor of tumor necrosis factor alpha-induced apoptosis.  
Li Y, Kang J, Friedman J, Tarassishin L, Ye J, Kovalenko A, Wallach D, Horwitz MS.  
Proc Natl Acad Sci U S A. 1999 Feb 2;96(3):1042-7.  
PMID: 9927690 [PubMed - indexed for MEDLINE]  
[Free article](#)
5. Epstein-Barr virus-transforming protein latent infection membrane protein 1 activates transcription factor NF-kappaB through a pathway that includes the NF-kappaB-inducing kinase and the IkappaB kinases IKKalpha and IKKbeta.  
Sylla BS, Hung SC, Davidson DM, Hatzivassiliou E, Malinin NL, Wallach D, Gilmore TD, Kieff E, Mosialos G.  
Proc Natl Acad Sci U S A. 1998 Aug 18;95(17):10106-11.  
PMID: 9707698 [PubMed - indexed for MEDLINE]  
[Free article](#)
6. CD27, a member of the tumor necrosis factor receptor superfamily, activates NF-kappaB and stress-activated protein kinase/c-Jun N-terminal kinase via TRAF2, TRAF5, and NF-kappaB-inducing kinase.  
Akiba H, Nakano H, Nishinaka S, Shindo M, Kobata T, Atsuta M, Morimoto C, Ware CF, Malinin NL, Wallach D, Yagita H, Okumura K.  
J Biol Chem. 1998 May 22;273(21):13353-8.  
PMID: 9582383 [PubMed - indexed for MEDLINE]  
[Free article](#)
7. TRAF2 plays a dual role in NF-kappaB-dependent gene activation by mediating the TNF-induced activation of p38 MAPK and IkappaB kinase pathways.  
Carpentier I, Declercq W, Malinin NL, Wallach D, Fiers W, Beyaert R.  
FEBS Lett. 1998 Mar 27;425(2):195-8.  
PMID: 9559646 [PubMed - indexed for MEDLINE]

8. MAP3K-related kinase involved in NF-kappaB induction by TNF, CD95 and IL-1.  
Malinin NL, Boldin MP, Kovalenko AV, Wallach D.  
*Nature*. 1997 Feb 6;385(6616):540-4.  
PMID: 9020361 [PubMed - indexed for MEDLINE]